

# Your personalized savings and support program\*

Scroll down for instructions on how to access a savings card.



Patient portrayals.

\*Only commercially insured patients are eligible for the TRINTELLIX savings card. The TRINTELLIX savings card cannot be used by patients in federal, state, or government-funded healthcare programs or by cash patients. See savings card for Eligibility Requirements and Terms & Conditions.

## WHAT IS TRINTELLIX?

TRINTELLIX is a prescription medicine used in adults to treat a certain type of depression called Major Depressive Disorder (MDD).

## IMPORTANT SAFETY INFORMATION

### **Suicidal Thoughts & Actions**

- **TRINTELLIX and other antidepressants may increase suicidal thoughts and actions in some people 24 years of age and younger.**
- **TRINTELLIX is not for use in children under 18.**
- **Call your doctor or get emergency help right away if you have new or sudden changes in mood, behavior, thoughts or feelings, or symptoms that are new, worse or worry you; or if you develop suicidal thoughts or actions.**

*Please see additional Important Safety Information on pages 3, 4, and 6 and accompanying Medication Guide, including Full Boxed WARNING for Suicidal Thoughts and Actions, on pages 7-8, and discuss with your doctor.*

Enroll now to receive savings\* and support at no cost to you.

A support program built to partner with you as you start your treatment with TRINTELLIX (vortioxetine).

**Customize your journey with support tools like:**



Savings card\*

\*Must meet eligibility requirements.

Pay as little as \$10 for a 30- or 90-day prescription. Maximum savings of \$100 on a 30-day or \$300 on a 90-day script.



Mobile wallet\*

\*Must meet eligibility requirements.

Download your savings card and add it to your mobile wallet for your next refill.



Informative emails

Receive emails with tools to help empower you to have meaningful discussions with your doctor.



Inspiring texts and refill reminders when needed

Uplifting messages and medication refill reminders sent to help you along the way.



Chats with a tAccess advisor

Call to receive information about the savings card and other helpful support features (1-844-654-TRIN).  
8:00 AM – 8:00 PM EST M-F

From savings\* to personalized support, tAccess is here to help.

**Please see additional Important Safety Information on pages 3, 4, and 6 and accompanying Medication Guide, including Full Boxed WARNING for Suicidal Thoughts and Actions, on pages 7-8, and discuss with your doctor.**

## IMPORTANT SAFETY INFORMATION (cont'd)

### **Suicidal Thoughts & Actions**

TRINTELLIX and other antidepressants may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. TRINTELLIX is not for use in children under 18. Depression or other mental illnesses are the most important causes of suicidal thoughts or actions. Pay close attention to any changes, especially new or sudden changes in mood, behavior, thoughts, or feelings. Call your doctor or get emergency help right away if you have symptoms such as suicidal thoughts or actions, impulsivity, aggressive or violent actions, depression, anxiety or panic attacks, agitation, restlessness, anger, irritability, trouble sleeping, an increase in activity or talking, or other unusual changes in behavior or mood; especially if they are new, worse, or worry you.

### **Who should not take TRINTELLIX?**

#### **Do not start or take TRINTELLIX if you:**

- are allergic to vortioxetine or any of the ingredients in TRINTELLIX
- take a Monoamine Oxidase Inhibitor (MAOI)
- have stopped taking an MAOI in the last 14 days
- are being treated with the antibiotic linezolid or intravenous methylene blue

Do not start taking an MAOI for at least 21 days after you stop treatment with TRINTELLIX.

### **What should I tell my doctor before taking TRINTELLIX?**

#### **Before taking TRINTELLIX, tell your doctor:**

- about all your medical and other health conditions
- if you are pregnant or plan to become pregnant, since TRINTELLIX may harm your unborn baby. Taking TRINTELLIX during your third trimester may cause your baby to have withdrawal symptoms after birth or to be at increased risk for a serious lung problem at birth. Tell your doctor right away if you become or think you are pregnant while taking TRINTELLIX.
- if you are breastfeeding or plan to breastfeed, since it is not known if TRINTELLIX passes into your breast milk



**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements, since TRINTELLIX and some medicines may cause serious side effects (or may not work as well) when taken together. **Especially tell your doctor if you take:** medicines for migraine headache called triptans; tricyclic antidepressants; opioids (such as fentanyl and tramadol); lithium; tryptophan; buspirone; St. John's Wort; medicines that can affect blood clotting such as aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), warfarin; and diuretics.

*Please see additional Important Safety Information on pages 4 and 6 and accompanying Medication Guide, including Full Boxed WARNING for Suicidal Thoughts and Actions, on pages 7-8, and discuss with your doctor.*

Pay as little as \$10 for a 30- or 90-day prescription.\*  
Ask your doctor what's right for you.

## HOW DO I GET STARTED?

To start saving\*

- 1 SIGN UP for the tAccess Support Program to get your savings card\*
- 2 ADD your savings card to a mobile wallet for future use! (Accessible via  )
- 3 USE your card to save on the next refill of your TRINTELLIX prescription

\*Commercially insured patients only. Must meet eligibility requirements.  
Maximum savings of \$100 on a 30-day or \$300 on a 90-day prescription.

Patients can join the tAccess Support Program.†

†Only commercially insured patients are eligible for the TRINTELLIX savings card.

## IMPORTANT SAFETY INFORMATION (cont'd)

### What are the possible side effects of TRINTELLIX?

**TRINTELLIX may cause serious side effects, including:**

- **Serotonin syndrome:** A potentially life-threatening problem that can happen when you take TRINTELLIX with certain other medicines. Call your doctor or go to the nearest emergency room right away if you have any of the following signs and symptoms of serotonin syndrome: agitation; seeing or hearing things that are not real; confusion; coma; fast heart beat; changes in blood pressure; dizziness; sweating; flushing; high body temperature; shaking, stiff muscles, or muscle twitching; loss of coordination; seizures; nausea, vomiting, diarrhea.
- **Increased risk of bleeding:** Taking TRINTELLIX with aspirin, NSAIDs, warfarin or blood thinners may add to this risk. Tell your doctor right away about any unusual bleeding or bruising.
- **Manic episodes:** Manic episodes may happen in people with bipolar disorder who take TRINTELLIX. Symptoms may include: greatly increased energy; racing thoughts; unusually grand ideas; talking more or faster than usual; severe problems sleeping; reckless behavior; excessive happiness or irritability.

*Please see additional Important Safety Information on pages 3 and 6 and accompanying Medication Guide, including Full Boxed WARNING for Suicidal Thoughts and Actions, on pages 7-8, and discuss with your doctor.*

# Start saving on your prescription today!

While all TRINTELLIX patients can join the tAccess Support Program, **only commercially insured patients are eligible to use a TRINTELLIX Savings Card.\***

\*All TRINTELLIX patients can join the tAccess Support Program. Only commercially insured patients are eligible for the TRINTELLIX savings card. See savings card for full Eligibility Requirements and Terms & Conditions.

**Find out if you're eligible to start saving.**

*Please see additional Important Safety Information on pages 3, 4, and 6 and accompanying Medication Guide, including Full Boxed WARNING for Suicidal Thoughts and Actions, on pages 7-8, and discuss with your doctor.*

## IMPORTANT SAFETY INFORMATION (cont'd)

### What are the possible side effects of TRINTELLIX? (cont'd)

- **Discontinuation syndrome:** Suddenly stopping TRINTELLIX may cause you to have serious side effects including: nausea; sweating; changes in your mood; irritability and agitation; dizziness; electric shock feeling; tremor; anxiety; confusion; headache; tiredness; problems sleeping; hypomania; ringing in your ears; seizures.
- **Eye problems:** TRINTELLIX may cause a type of eye problem called angle-closure glaucoma in people with certain other eye conditions. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. Call your doctor if you have eye pain, changes in your vision, or swelling or redness in or around the eye.
- **Low levels of salt (sodium) in your blood:** Low sodium levels in your blood that may be serious and may cause death can happen during treatment with TRINTELLIX. Elderly people and people who take certain medicines may be at a greater risk for developing low sodium levels in your blood. Signs and symptoms may include headache; difficulty concentrating; memory changes; confusion; weakness and unsteadiness on your feet which can lead to falls. **In more severe or more sudden cases, signs and symptoms include:** seeing or hearing things that are not real; fainting; seizures; coma; stopping breathing.
- **Sexual problems:** Taking antidepressants like TRINTELLIX may cause sexual problems. Symptoms in males may include: delayed ejaculation or inability to have an ejaculation, decreased sex drive, or problems getting or keeping an erection. Symptoms in females may include: decreased sex drive, or delayed orgasm or inability to have an orgasm. Talk to your healthcare provider if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with TRINTELLIX.

### **The most common side effects of TRINTELLIX include:**

- nausea
- constipation
- vomiting

These are not all the possible side effects of TRINTELLIX. Tell your doctor if you have any side effect that bothers you or does not go away.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

***For additional Important Safety Information, click here for [Medication Guide](#), or see pages 7-8 of this brochure, and discuss with your doctor.***

TRINTELLIX is a trademark of H. Lundbeck A/S registered with the U.S. Patent and Trademark Office and used under license by Takeda Pharmaceuticals America, Inc. TAKEDA and the TAKEDA logo are registered trademarks of Takeda Pharmaceutical Company Limited. ©2022 Takeda Pharmaceuticals U.S.A., Inc. All rights reserved. 1-877-TAKEDA-7 (1-877-825-3327).  
US-VOR-0908v1.0 1/22



## MEDICATION GUIDE

TRINTELLIX® (trin'-TELL-ix)

(vortioxetine)

Tablets

### What is the most important information I should know about TRINTELLIX?

#### TRINTELLIX can cause serious side effects, including:

• **Increased risk of suicidal thoughts and actions.** TRINTELLIX and other antidepressant medicines may increase suicidal thoughts and actions in some people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. TRINTELLIX is not for use in children.

• **Depression or other mental illnesses are the most important causes of suicidal thoughts or actions.**

#### How can I watch for and try to prevent suicidal thoughts and actions?

- Pay close attention to any changes, especially sudden changes in mood, behavior, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed.
- Call your healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings or if you develop suicidal thoughts or actions.
- Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms.

**Call your healthcare provider or get emergency help right away if you have any of the following symptoms, especially if they are new, worse, or worry you:**

- attempts to commit suicide
- acting on dangerous impulses
- acting aggressive or violent
- thoughts about suicide or dying
- new or worse depression
- new or worse anxiety or panic attacks
- feeling agitated, restless, angry or irritable
- trouble sleeping
- an increase in activity or talking more than what is normal for you
- other unusual changes in behavior or mood

### What is TRINTELLIX?

TRINTELLIX is a prescription medicine used in adults to treat a certain type of depression called Major Depressive Disorder (MDD).

#### Do not take TRINTELLIX if you:

- are allergic to vortioxetine or any of the ingredients in TRINTELLIX. See the end of this Medication Guide for a complete list of ingredients in TRINTELLIX
- take a Monoamine Oxidase Inhibitor (MAOI)
- have stopped taking an MAOI in the last 14 days
- are being treated with the antibiotic linezolid or intravenous methylene blue

Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI or one of these medicines, including the antibiotic linezolid or intravenous methylene blue.

**Do not start TRINTELLIX if you stopped taking an MAOI in the last 14 days.**

**Do not start taking an MAOI for at least 21 days after you stop treatment with TRINTELLIX.**

#### Before taking TRINTELLIX, tell your healthcare provider about all your medical conditions, including if you:

- have or had seizures or convulsions
- have or had bleeding problems

- have, or have a family history of, bipolar disorder, mania or hypomania
- have high pressure in the eye (glaucoma)
- have low sodium levels in your blood
- are pregnant or plan to become pregnant. TRINTELLIX may harm your unborn baby. Taking TRINTELLIX during your third trimester of pregnancy may cause your baby to have withdrawal symptoms after birth or may cause your baby to be at increased risk for a serious lung problem at birth. Talk to your healthcare provider about the risks to you and your unborn or newborn baby if you take TRINTELLIX during pregnancy.
  - Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with TRINTELLIX.
  - There is a pregnancy registry for females who are exposed to TRINTELLIX during pregnancy. The purpose of the registry is to collect information about the health of females exposed to TRINTELLIX and their baby. If you become pregnant during treatment with TRINTELLIX, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants at 1-844-405-6185 or visit online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/>.
- are breastfeeding or plan to breastfeed. It is not known if TRINTELLIX passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with TRINTELLIX.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

TRINTELLIX and some other medicines may affect each other causing possible serious side effects. TRINTELLIX may affect the way other medicines work and other medicines may affect the way TRINTELLIX works.

#### Especially tell your healthcare provider if you take:

- medicines used to treat migraine headache called triptans
- tricyclic antidepressants
- fentanyl
- lithium
- tramadol
- tryptophan
- buspirone
- St. John's Wort
- medicines that can affect blood clotting such as aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), warfarin
- diuretics

Ask your healthcare provider if you are not sure if you are taking any of these medicines. Your healthcare provider can tell you if it is safe to take TRINTELLIX with your other medicines.

Do not start or stop any other medicines during treatment with TRINTELLIX without talking to your healthcare provider first. Stopping TRINTELLIX suddenly may cause you to have serious side effects. See, "What are the possible side effects of TRINTELLIX?"

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

#### How should I take TRINTELLIX?

- Take TRINTELLIX exactly as your healthcare provider tells you to take it. Your healthcare provider may need to change the dose of TRINTELLIX until it is the right dose for you.
- Take TRINTELLIX 1 time each day.
- TRINTELLIX may be taken with or without food.
- If you take too much TRINTELLIX, call your poison control center at 1-800-222-1222 or go to the nearest hospital emergency room right away.

## What are the possible side effects of TRINTELLIX?

### TRINTELLIX may cause serious side effects, including:

- See, “What is the most important information I should know about TRINTELLIX?”
- **Serotonin syndrome.** A potentially life-threatening problem called serotonin syndrome can happen when you take TRINTELLIX with certain other medicines. See, “Who should not take TRINTELLIX?”  
**Call your healthcare provider or go to the nearest hospital emergency room right away** if you have any of the following signs and symptoms of serotonin syndrome:
  - agitation
  - sweating
  - seeing or hearing things that are not real (hallucinations)
  - flushing
  - confusion
  - high body temperature (hyperthermia)
  - coma
  - shaking (tremors), stiff muscles, or muscle twitching
  - fast heart beat
  - loss of coordination
  - changes in blood pressure
  - seizures
  - dizziness
  - nausea, vomiting, diarrhea
- **Increased risk of bleeding.** Taking TRINTELLIX with aspirin, non-steroidal anti-inflammatory drugs (NSAIDs), warfarin or blood thinners may add to this risk. Tell your healthcare provider right away about any unusual bleeding or bruising.
- **Manic episodes.** Manic episodes may happen in people with bipolar disorder who take TRINTELLIX. Symptoms may include:
  - greatly increased energy
  - severe problems sleeping
  - racing thoughts
  - reckless behavior
  - unusually grand ideas
  - excessive happiness or irritability
  - talking more or faster than usual
- **Discontinuation syndrome.** Suddenly stopping TRINTELLIX may cause you to have serious side effects. Your healthcare provider may want to decrease your dose slowly. Symptoms may include:
  - nausea
  - anxiety
  - electric shock feeling (paresthesia)
  - hypomania
  - irritability and agitation
  - tiredness
  - confusion
  - sweating
  - ringing in your ears (tinnitus)
  - tremor
  - dizziness
  - problems sleeping
  - headache
  - changes in your mood
  - seizures
- **Eye problems (angle-closure glaucoma).** TRINTELLIX may cause a type of eye problem called angle-closure glaucoma in people with certain other eye conditions. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. Call your healthcare provider if you have eye pain, changes in your vision, or swelling or redness in or around the eye.
- **Low levels of sodium in your blood (hyponatremia).** Low sodium levels in your blood that may be serious and may cause death, can happen during treatment with TRINTELLIX. Elderly people and people who take certain medicines may be at a greater risk for developing low sodium levels in your blood. Signs and symptoms may include:
  - headache
  - difficulty concentrating
  - memory changes
  - confusion
  - weakness and unsteadiness on your feet which can lead to falls

### In more severe or more sudden cases, signs and symptoms include:

- seeing or hearing things that are not real (hallucinations)
- fainting
- seizures
- coma
- stopping breathing (respiratory arrest)
- **Sexual problems (dysfunction).** Taking TRINTELLIX may cause sexual problems.  
Symptoms in males may include:
  - Delayed ejaculation or inability to have an ejaculation
  - Decreased sex drive
  - Problems getting or keeping an erectionSymptoms in females may include:
  - Decreased sex drive
  - Delayed orgasm or inability to have an orgasm

Talk to your healthcare provider if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with TRINTELLIX. There may be treatments your healthcare provider can suggest.

**The most common side effects of TRINTELLIX include** nausea, constipation, vomiting.

These are not all the possible side effects of TRINTELLIX.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### How should I store TRINTELLIX?

- Store TRINTELLIX at room temperature between 68°F to 77°F (20°C to 25°C).
- **Keep TRINTELLIX and all medicines out of the reach of children.**

### General information about the safe and effective use of TRINTELLIX.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not take TRINTELLIX for a condition for which it was not prescribed. Do not give TRINTELLIX to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about TRINTELLIX that is written for healthcare professionals.

### What are the ingredients in TRINTELLIX?

**Active ingredient:** vortioxetine hydrobromide

**Inactive ingredients:** mannitol, microcrystalline cellulose, hydroxypropyl cellulose, sodium starch glycolate, magnesium stearate and film coating consisting of hypromellose, titanium dioxide, polyethylene glycol 400, iron oxide red (5 mg and 20 mg) and iron oxide yellow (10 mg)

Distributed and Marketed by:

**Takeda Pharmaceuticals America, Inc.** Lexington, MA 02421

Marketed by:

**Lundbeck,** Deerfield, IL 60015

TRINTELLIX is a trademark of H. Lundbeck A/S registered with the U.S. Patent and Trademark Office and used under license by Takeda Pharmaceuticals America, Inc.

All other trademarks are the property of their respective owners.

©2013-2021 Takeda Pharmaceuticals America, Inc.

For more information, go to [www.TRINTELLIX.com](http://www.TRINTELLIX.com) or call 1-877-TAKEDA-7 (1-877-825-3327).

This Medication Guide has been approved by the U.S. Food and Drug Administration

Revised: 09/2021

SPI-0334